各都道府県衛生主管部(局)薬務主管課 御中

厚生労働省医薬・生活衛生局医薬品審査管理課

第十八改正日本薬局方(英文版)正誤表の送付について(その2)

第十八改正日本薬局方(令和3年厚生労働省告示第220号)の英文版につきまして、 一部に誤植等がありましたので別紙のとおり正誤表を送付いたします。 Bicalutamide ビカルタミド

Page	Line	Correction	Error
		Determine each peak area by the automatic	Determine each peak area by the automatic
		integration method: the peak areas of related	integration method: the peak areas of related
		substance M, having the relative retention time	substance M, having the relative retention time
		of about 0.26 to bicalutamide, related	of about 0.26 to bicalutamide, related
		substance N, having the relative retention time	substance N, having the relative retention time
550	left 21-20	of about 0.34, related substance K, having the	of about 0.34, related substance L, having the
550	lett 21-20	relative retention time of about 1.03 and	relative retention time of about 1.03 and
		related substance L, having the relative	related substance K, having the relative
		retention time of about 1.13, obtained from the	retention time of about 1.13, obtained from the
		sample solution, are not larger than the peak	sample solution, are not larger than the peak
		area of bicalutamide from the standard	area of bicalutamide from the standard
		solution,	solution,

Candesartan Cilexetil and Amlodipine Besylate Tablets カンデサルタンシレキセチル・アムロジピンベシル酸塩錠

Page	Line	Correction	Error
615-618		Amlodipine Bes <u>i</u> late	Amlodipine Bes <u>v</u> late

Imidapril Hydrochloride Tablets イミダプリル塩酸塩錠

	Page	Line	Correction	Error
Ī	1143	left 29-28	Add diluted methanol (2 in 5) to make 50 mL,	Add diluted ethanol (2 in 5) to make 50 mL,

Zopiclone ゾピクロン

Page	Line	Correction	Error
1935	right 33-36	determine each peak area by the automatic integration method: the peak areas of related substance A, having the relative retention time of about 0.1 to zopiclone, related substance B, having the relative retention time of about 0.2, related substance C, having the relative retention time of about 0.5, related substance D, having the relative retention time of about 0.9, obtained from the sample solution are not larger than 1/10 times the peak area of zopiclone from the standard solution, and the area of the peak other than zopiclone and the peaks mentioned above from the sample solution is not lager than 1/10 times the peak area of zopiclone from the standard solution.	determine each peak area by the automatic integration method: the peak areas of related substance A, having the relative retention time of about 0.1 to zopiclone, related substance B, having the relative retention time of about 0.2, related substance C, having the relative retention time of about 0.5, related substance D, having the relative retention time of about 0.9 and the peaks other than mentioned above, obtained from the sample solution, are not larger than 1/10 times the peak area of zopiclone from the standard solution.